

SOL-TRACE[®]

SOLUBLE TRACE ELEMENT MIX

For supplementing and overcoming trace element deficiency

GUARANTEED ANALYSIS

Sulfur (S)	14.00%
14.00% Combined Sulfur (S)	
Boron (B)	1.45%
Copper (Cu).....	3.20%
3.20% Water Soluble Copper (Cu)	
Iron (Fe).....	7.50%
7.50% Water Soluble Iron (Fe)	
Manganese (Mn)	8.15%
8.15% Water Soluble Manganese (Mn)	
Molybdenum (Mo).....	0.046%
Zinc (Zn)	4.50%
4.50% Water Soluble Zinc (Zn)	

Derived from Borax, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Sodium Molybdate and Zinc Sulfate. F593

Information regarding the content and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

CAUTION: Contains Boron. The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crop. **WARNING:** The use of fertilizing material containing molybdenum (Mo) may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

SOIL MUST BE WET BEFORE APPLYING OR PLANT INJURY CAN OCCUR.

RATES OF APPLICATION FOR NURSERY & GREENHOUSE USE

This material is immediately available when applied and should show results within days. Do not over apply. Use soil and tissue test as a guide. When on a constant feeding schedule, using basic NPK fertilizers, the normal rate of application is 2 ozs. of SOL-TRACE for every 25 lbs. (5 grams per kg) of fertilizer material being applied. Not recommended for use with trace element fortified fertilizers. As an occasional or supplemental application to bench crops, use 2 ozs. per 100 sq. ft. (6.10 grams per square meter). Do not repeat unless deficiency has definitely been established. For potted crops, use 2 ozs. in 25 gals. (15 grams per 25 liters). Apply as a normal saturation of root area. No repeat should be necessary.

TURF - GREENS, TEES & FAIRWAYS

For one time corrective action - Use 1/4 to 1/2 oz. per 1000 square feet in 5 to 25 gallons of water

WARNING! AVOID CONTACT WITH EYES AND SKIN TO AVOID IRRITATION. WASH CLOTHING THOROUGHLY BEFORE RE-USE.

- FIRST AID -

IN CASE OF CONTACT, IMMEDIATELY FLUSH SKIN OR EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. IF IRRITATION PERSISTS, CALL PHYSICIAN. FOR PROFESSIONAL USE ONLY.